

# PARABOLAN L.A

## TREMBOLONE ENANTHATE 200 mg/ml. MUSCLE INJECTABLE OIL SOLUTION/INTRAMUSCULAR WAY

### Composition of this medicine

Every 1 ml. of Injectable Oily Solution contains:

Trenbolone enanthate..... 200 mg.  
Excipientes..... c.s.p.

**What is Trenbolone?** It is a medicine that contains Trenbolone Enanthate which belongs to the group of Androgens. Trenbolone is basically in charge of developing and maintaining the body in recovery from debilitating catabolic processes and helping various systems, including the muscle.

**How does this medicine work?** It works by activating its very selective anti-catabolic properties, helping to maintain protein values in the body, increasing erythropoiesis and raising the state of conservation and development of the muscular system, as well as regulating fats.

**Who should be careful when using this medication?** Those with heart disease, high blood pressure, tumors, breast and prostate cancer, prostatic hyperplasia, behavioral disorders, liver problems, poor kidney function, imbalanced calcium in the blood, thyroid, acne, excessively oily skin, pictures severe or chronic respiratory Especially in women, due to its hormonal action, it is prohibited and contraindicated. Also children, adolescents and seniors. Pregnant and lactating women as it is potentially risky. It should not be administered to those who receive other immunodebitating treatments, they must necessarily consult their doctor.

**THERAPEUTIC ACTION** For the treatment of diseases and conditions that require an increase in protein synthesis.

### POSOLOGY

Recommended dose in men 150 mg per week, or according to medical criteria.

**ACTION MECHANISMS** Trenbolone is a substance that belongs to the group of non-stilbene xenobiotics, therefore it is a synthetic hormone derived from Nortestosterone, thus being a modified derivative of Nandrolone and differs from it by including two bonds with double characteristics. in Carbons 9 and 11, which inhibits its aromatization so characteristic of this type of hormones (9-en), increasing its androgenic affinity and slowing down its metabolism. At the C9 and C10 bond level, it inhibits Estrogenic activity, although it seems that this substance itself is not metabolically canceled since these compounds are necessary for it to become Estrogen, so the final steroid (Trenbolone) is 3 times more powerful in its action anabolic as androgenic than nandrolone base. Since Trenbolone is not aromatized in the body, it is no longer Estrogenic, presenting a greater affinity for Progesterone receptors, therefore the benefits are demonstrated in that the side effects due to the aromatization of Estrogen are practically not present and because the already modified Trenbolone continues acting as a highly potent muscle mass, strength and definition drug, much more so than Testosterone. But the most striking thing is that due to its steroidal characteristics, Trenbolone reaches a greater amount of time within the body after its administration, in addition to the strong impact it has on endogenous testosterone, it is safer to use drugs such as HCG after finishing a treatment cycle since without the use of this type of drugs, the hormonal leveling of endogenous gonadotrophins at the testicular level could last for a long time

**ADVERSE BIOLOGICAL EFFECTS OF TREMBOLONE IN GENERAL** Woman: Severe virilization, Larynx: Growth and elongation of the vocal cords, Skin: Stimulates the production of fat and Acne, Prostate: Stimulates its growth (HBP), Brain: Temporary inhibition of respiratory centers especially at the time of administration (Tren Cough ), Sexual and reproductive disorders due to high levels of Prolactin, Increased aggressiveness (rare)

**CONTRAINDICATIONS:** Cardiovascular Diseases, Arterial Hypertension, Thyroid and Adrenal Diseases. Neurological and Prostatic Diseases. (Cancer) Active Breast Cancer, Kidney Diseases, Liver Diseases of any etiology, Pregnancy, Breastfeeding, COPD, EBOC, Personality Disorders (aggressiveness).

**WARNINGS and CAUTIONS:** It should not be used for long periods. Do not administer to patients with other treatments. Do not consume alcohol simultaneously. Do not exceed doses. Keep out of the reach of children. Administer with caution to patients who are allergic or sensitive to this medicine or its components. **ATTENTION!** Warnings for athletes have NOT been described, but it is reported that this medicine contains a component that in a drug test can give the result as POSITIVE (+).

**ADVERSE REACTIONS:** The appearance of adverse reactions can be; Virilization in women (increased tone of voice, hirsutism, clitoral hypertrophy, menstrual irregularities, amenorrhea, thickening of the skin and others) In adults Acne, moderate polycythemia and obstruction due to Prostatic Hyperplasia, Liver Diseases (Cholestatic Jaundice), Hepatic Carcinoma, Mild Hyposaline Retention Prostate Cancer, Arterial Hypertension. In small men and mild feminizing effects (mild gynecomastia, decreased libido, probable erection disorders and androgenic-type impotence), especially at the end of the administration cycle. Hormonal alterations in Thyroid. Nephropathies and alteration of the Adrenals.

**USE RESTRICTIONS:** En especial en las mujeres por su acción hormonal se encuentra prohibida y contraindicada. También los niños, adolescentes y personas de la tercera edad. Embarazadas y las que dan de lactar ya que es potencialmente riesgoso.

### OVERDOSE:

In case of an overdose, quickly contact the Toxicology Center of the Hospital for Medical Emergencies in Paraguay

### PRESENTATION

Box containing a vial of 10 mL  
Box containing 5 vials of 1 mL  
Box containing 10 vials of 1 mL

Store at room temperature between 15°C - 30°C.

Keep out of the reach of children.